

Product datasheet for **TA396645S**

ACHE Goat Polyclonal Antibody

Product data:

| | |
|-----------------------|--|
| Product Type: | Primary Antibodies |
| Applications: | ELISA, IP, WB |
| Recommended Dilution: | WB: 1:500 - 1:5,000 ELISA: 1:5,000 - 1:20,000 |
| Host: | Goat |
| Clonality: | Polyclonal |
| Immunogen: | Acetyl Cholinesterase [Bovine Erythrocytes] |
| Specificity: | Anti-Acetyl Cholinesterase is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum as well as purified and partially purified Acetyl Cholinesterase [Bovine Erythrocytes]. Cross reactivity against Acetyl Cholinesterase from other sources is unknown. |
| Formulation: | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Concentration: | 1.0 mg/ml - lot specific |
| Conjugation: | Unconjugated |
| Storage: | Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing. |
| Stability: | Expiration date is one (1) year from date of receipt. |
| Database Link: | P23795 |
| Background: | Acetylcholinesterase terminates signal transduction at the neuromuscular junction by rapid hydrolysis of the acetylcholine released into the synaptic cleft. |
| Synonyms: | goat anti-Acetyl Cholinesterase Antibody, Acetylcholinesterase, AChE |



[View online »](#)

Note: Anti-Acetyl Cholinesterase Antibody has been tested by western blot and is suitable to be assayed against 1.0 µg of Acetyl Cholinesterase [Bovine Erythrocytes] in a standard ELISA using Peroxidase conjugated Affinity Purified anti-Goat IgG [H&L] (Rabbit) code #605-4302 and (ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:1,200 to 1:5,000 of the reconstitution concentration is suggested for this product.