

## Product datasheet for **TA396662S**

### **FOLR1 Goat Polyclonal Antibody**

#### **Product data:**

Product Type:	Primary Antibodies
Applications:	ELISA, IP, WB
Recommended Dilution:	<b>WB:</b> 1:500 - 1:1,000 <b>ELISA:</b> 1:1,000 - 1:2,500
Host:	Goat
Clonality:	Polyclonal
Immunogen:	Folate Binding Protein [Bovine Milk]
Specificity:	Anti-Folate-binding protein is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum as well as purified and partially purified Folate Binding Protein [Bovine Milk]. Cross reactivity against Folate Binding Protein from other sources is unknown.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Concentration:	1.0 mg/mL - lot specific
Conjugation:	HRP
Storage:	Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.
Stability:	Expiration date is one (1) year from date of receipt.
Database Link:	<u><a href="#">P02702</a></u>



[View online »](#)

<b>Background:</b>	Folate-binding protein (FBP) are proteins that bind folate, typically the folate receptors and reduced folate carrier. In cancer research it can be a marker for ovarian cancer. It can be measured to help in tumor classification to aid treatment. FBP is over-expressed (20-80 fold) in more than 90% of ovarian and endometrial cancers, as well as 20-50% of breast, lung, colorectal, and renal cell carcinomas. FBP has very limited tissue distribution and expression in non-malignant tissue, making it a good immunotherapy target.
<b>Synonyms:</b>	goat anti-Folate Binding Protein Antibody HRP Conjugation, Peroxidase Conjugated goat anti-Folate Binding Protein Antibody, Folate receptor alpha, FR-alpha, Folate receptor 1, Milk folate-binding protein, Folate-binding protein 1, FBP
<b>Note:</b>	Anti-Folate Binding Protein has been tested in western blot and is suitable to be assayed against 1.0 ug of Folate Binding Protein [Bovine Milk] in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:2,000 to 1:10,000 of the reconstitution concentration is suggested for this product.