

Product datasheet for **TA396672S**

NAM-DH Goat Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, IP, WB
Recommended Dilution:	WB: 1:500 - 1:5,000 ELISA: 1:5,000 - 1:20,000
Reactivity:	Bacterial
Host:	Goat
Clonality:	Polyclonal
Immunogen:	N-Acyl Mannosamine-1-Dehydrogenase [Recombinant Sequence from Pseudomonas expressed in E.coli]
Specificity:	N-Acylmannosamine-1-Dehydrogenase is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, and purified and partially purified N-Acylmannosamine-1-Dehydrogenase derived from Pseudomonas.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Concentration:	1 mg/mL - lot specific
Conjugation:	HRP
Storage:	Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.
Stability:	Expiration date is one (1) year from date of receipt.
Database Link:	P22441
Background:	N-Acylmannosamine-1-Dehydrogenase acts on acetyl-D-mannosamine and glycolyl-D-mannosamine. This enzyme belongs to the family of oxidoreductases, specifically those acting on the CH-OH group of donor with NAD ⁺ or NADP ⁺ as acceptor.




[View online »](#)

Synonyms: goat anti-N-Acylmannosamine-1-Dehydrogenase Antibody HRP Conjugation, Peroxidase Conjugated goat anti-N-Acylmannosamine-1-Dehydrogenase Antibody, N-acylmannosamine 1-dehydrogenase, NAM-DH

Note: Anti-N-Acylmannosamine-1-Dehydrogenase has been assayed against 1.0 ug of N-Acylmannosamine-1-Dehydrogenase IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:4,000 to 1:16,000 is suggested for this product.

Product images:



Rockland Goat anti N-acylmannosamino-1-Dehydrogenase antibody (200-101-228 lot 8181) was used to detect purified N-acylmannosamino-1-Dehydrogenase under reducing (R) and non-reducing (NR) conditions. Reduced samples of purified target proteins contained 4% BME and were boiled for 5 minutes. Samples of ~1ug of protein per lane were run by SDS-PAGE. Protein was transferred to nitrocellulose and probed with 1:3000 dilution of primary antibody (ON 4 C in MB-070). Detection shown was using Dylight 488 conjugated Donkey anti goat (605-741-125 lot 21094 1:10K in TBS/MB-070 1 hr RT). Images were collected using the BioRad VersaDoc System.