

Product datasheet for **TA396687**

Goat Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	WB: 1:1,000 - 1:5,000 ELISA: 1:10,000 - 1:25,000
Host:	Goat
Clonality:	Polyclonal
Immunogen:	Luciferase [Photinus pyralis (Firefly)]
Specificity:	Anti-Luciferase Biotin Conjugated is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Goat Serum as well as purified and partially purified Luciferase [Photinus pyralis (Firefly)]. No reactivity is observed against Sea pansy (<i>Renilla reniformis</i>) luciferase.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reconstitution Method:	Restore with deionized water (or equivalent) - Reconstitution Volume: 1.0 mL
Concentration:	10.0 mg/mL - lot specific
Conjugation:	Biotin
Storage:	Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.
Stability:	Expiration date is one (1) year from date of receipt.



[View online »](#)

- Background:** Anti luciferase Antibody recognizes luciferase that is commonly used in biological research as a reporter to assess the transcriptional activity in cells that are transfected with a genetic construct containing the luciferase gene under the control of a promoter of interest. Luciferase can also be used to detect the level of cellular ATP in cell viability assays or for kinase activity assays. Additionally proluminescent molecules that are converted to luciferin upon activity of a particular enzyme can be used to detect enzyme activity in coupled or two-step luciferase assays. Such substrates have been used to detect caspase activity and cytochrome P450 activity, among others.
- Synonyms:** goat anti-Luciferase Antibody biotin Conjugation, Biotin Conjugated goat anti-Luciferase Antibody, Luciferase Biotin, Anti-Luciferase BAC Antibody
- Note:** Anti-Luciferase Peroxidase conjugated antibody is produced from Photinus pyralis (Firefly) which produces a green light with a wavelength of 562 nm. Anti-Luciferase Antibody has been tested by ELISA and western blot. Expect a band at approximately 60.7 kDa in size corresponding to Luciferase by western blotting in the appropriate cell lysate or extract. Anti-Luciferase has been assayed against 1.0 ug of Luciferase in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:80,000 to 1:310,000 of the reconstitution concentration is suggested for this product.