

Product datasheet for **TA396694S**

choA Goat Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, IP, WB
Recommended Dilution:	WB: 1:500 - 1:5,000 ELISA: 1:5,000 - 1:20,000
Reactivity:	Streptomyces
Host:	Goat
Clonality:	Polyclonal
Immunogen:	Cholesterol Oxidase [Streptomyces]
Specificity:	Anti-Cholesterol Oxidase is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Goat Serum as well as purified and partially purified Cholesterol Oxidase [Microorganism]. Cross reactivity against Cholesterol Oxidase from other sources is unknown.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Concentration:	1 mg/ml - lot specific
Conjugation:	Biotin
Storage:	Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.
Stability:	Expiration date is one (1) year from date of receipt.
Database Link:	P12676



[View online »](#)

Background: Anti-Cholesterol Oxidase antibody is a member of the GMC oxidoreductase family. Cholesterol Oxidase acts as a catalyst in the oxidation of the 3-beta-hydroxy group of cholesterol and the isomerization of the double bond of new product. Anti-Cholesterol Oxidase is ideal for investigators interested in Metabolism, Signal Transduction, or Cardiovascular research.

Synonyms: goat anti-Cholesterol Oxidase Antibody biotin Conjugation, biotin Conjugated goat anti-Cholesterol Oxidase Antibody, choA antibody, choD antibody, choM antibody, choS antibody, COD antibody

Note: Anti-Cholesterol Oxidase has been tested by ELISA, dot blot, and western blot. This product is assayed against 1.0 ug of Cholesterol Oxidase in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:4,000 to 1:20,000 of the reconstitution concentration is suggested for this product.

Product images:



Rockland Goat anti Cholesterol oxidase antibody (200-101-213 lot 6534) was used to detect purified cholesterol oxidase under reducing (R) and non-reducing (NR) conditions. Reduced samples of purified cholesterol oxidase contained 4% BME and were boiled for 5 minutes. Samples of ~1ug of protein per lane were run by SDS-PAGE. Protein was transferred to nitrocellulose and probed with a 1:3000 dilution of primary antibody (ON 4 C in MB-070). Detection shown was using Dylight 488 conjugated Donkey anti goat (605-741-125 lot 21094 1:10K in TBS/MB-070 1 hr RT). Images were collected using the BioRad VersaDoc System.