

## Product datasheet for **TA396720S**

### ALPI Goat Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, IHC, WB
Recommended Dilution:	<b>WB:</b> 1:1,000 - 1:5,000 <b>IHC:</b> User Optimized <b>ELISA:</b> 1:10,000 - 1:50,000
Host:	Goat
Clonality:	Polyclonal
Immunogen:	Alkaline Phosphatase [Calf Intestine]
Specificity:	Alkaline Phosphatase is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum as well as purified and partially purified Alkaline Phosphatase [Calf Intestine]. Cross reactivity against Alkaline Phosphatase from other tissues and species may occur but have not been specifically determined.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Concentration:	1.0 mg/mL - lot specific
Conjugation:	HRP
Storage:	Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.
Stability:	Expiration date is one (1) year from date of receipt.
Database Link:	<a href="#">P19111</a>



[View online »](#)

- Background:** Alkaline Phosphatase is present in all tissues throughout the entire body, but is particularly concentrated in liver, bile duct, kidney, bone, and the placenta. Humans and most other mammals contain the following alkaline phosphatase isozymes: ALPI – intestinal, ALPL – tissue non-specific (liver/bone/kidney), ALPP – placental (Regan isozyme). Elevated ALP indicates that there could be active bone formation occurring as ALP is a byproduct of osteoblast activity (such as the case in Paget's disease of bone). Levels are also elevated in people with untreated Coeliac Disease. All mammalian alkaline phosphatase isoenzymes except placental (PALP and SEAP) are inhibited by homoarginine, and, in similar manner, all except the intestinal and placental ones are blocked by levamisole. Heating for ~2 hours at 65°C inactivated most isoenzymes except Placental isoforms (PALP and SEAP).
- Synonyms:** goat anti-Alkaline Phosphatase Antibody HRP Conjugation, Peroxidase Conjugation goat anti-Alkaline Phosphatase Antibody, Alkaline phosphomonoesterase antibody, ALPI antibody, Glycerophosphatase antibody, IAP antibody, Intestinal alkaline phosphatase antibody, Kasahara isozyme antibody
- Note:** Anti-Alkaline Phosphatase peroxidase has been tested by dot blot and is functional in ELISA, Western blot, and Immunohistochemical assays. Optimization should be reached by researcher.