

Product datasheet for **TA396771S**

IDO1 Mouse Monoclonal Antibody [Clone ID: 10.1]

Product data:


Product Type:	Primary Antibodies
Clone Name:	10.1
Applications:	ELISA, IF, IHC, IP, WB
Recommended Dilution:	WB: 1:500-1:1500 IHC: 1:100-1:500 IF: 1:50-1:100 ELISA: 1:5000-1:50000
Reactivity:	Human, Mouse
Host:	Mouse
Isotype:	IgG3
Clonality:	Monoclonal
Immunogen:	Anti-IDO1 (MOUSE) Monoclonal Antibody was produced in mouse by repeated immunizations with fragment of recombinant human and mouse IDO1 protein followed by hybridoma development.
Specificity:	Anti-IDO1 was purified from concentrated tissue culture supernate by Protein G chromatography followed by extensive dialysis against the buffer stated above. This antibody is specific for human and mouse IDO1 protein. A BLAST analysis was used to suggest cross-reactivity with IDO1 from human and mouse sources based on 100% homology with the immunizing sequence. Cross-reactivity with IDO1 from other sources has not been determined.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Concentration:	1.0 mg/mL - lot specific
Conjugation:	Unconjugated
Storage:	Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.



[View online »](#)

Stability:	Expiration date is one (1) year from date of receipt.
Gene Name:	indoleamine 2,3-dioxygenase 1
Database Link:	Entrez Gene 3620 Human P14902
Background:	Indoleamine 2, 3-dioxygenase1 (IDO1) is a 41-42 kD intracellular enzyme that catabolizes tryptophan into kynurenine. IDO1 modulates levels of the amino acid tryptophan, which is vital for cell growth, but is also involved in the suppression of the immune response. IDO1 effects on immune suppression are due to decreased tryptophan availability and the generation of tryptophan metabolites, resulting in negative effects on T lymphocytes, including proliferation, function and survival. IDO1 may be involved in the suppression of the immune response to tumors, and blocking the IDO1 pathway may be a potential target for immuno and cancer therapy. IDO1 is expressed in a wide variety of tissues and can be upregulated by interferon gamma and other inflammatory cytokines.
Synonyms:	mouse anti-IDO1 antibody, Ido, Indo, Indoleamine 2,3-dioxygenase 1, Indoleamine-pyrrole 2,3-dioxygenase, Ido1, Ido-1
Note:	Anti-IDO1 antibody has been tested in ELISA, IP, and Western Blot. This antibody is suitable for use in IHC and Flow Cytometry. Specific conditions for reactivity should be optimized by the end user.

Product images:



Western Blot of Mouse Anti-IDO1 Antibody. Lane 1: untreated HeLa cells (p/n W09-000-364). Lane 2: IFN-r treated HeLa cells. Load: 35 µg per lane. Primary antibody: IDO 1 Antibody at 1:1000 for overnight at 4°C. Secondary antibody: IRDye800™ mouse secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO (p/n [B501-0500]) overnight at 4°C. Predicted/Observed size: 41-42 kDa, 41-42 kDa for IDO-1. Other band(s): none.