

Product datasheet for **TA396781**

GSTK1 Mouse Monoclonal Antibody [Clone ID: 16A11.E4.E3.E7]

Product data:

Product Type:	Primary Antibodies
Clone Name:	16A11.E4.E3.E7
Applications:	ELISA, IHC, IP, WB
Recommended Dilution:	WB: 1:500 - 1:1000 IHC: User Optimized ELISA: 1:50,000 - 1:200,000
Reactivity:	Human
Host:	Mouse
Isotype:	IgG2a, lambda
Clonality:	Monoclonal
Immunogen:	This Protein A purified antibody was prepared by repeated immunizations in mice with K1 Protein.
Specificity:	This product is purified from roller bottle culture by Protein A chromatography followed by extensive dialysis against the buffer stated above. Reacts specifically with K1 protein. Cross reactivity from other sources has not been determined.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Concentration:	1.0 mg/ml - lot specific
Conjugation:	Unconjugated
Storage:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Stability:	Expiration date is one (1) year from date of receipt.
Database Link:	Q9Y2Q3




[View online »](#)

Background: Rockland produces a wide range of human GST antibodies in our laboratories. Select appropriate GST antibodies for your research by isotype, epitope, applications and species reactivity. There are 22 members of the human GST family of proteins. GST is responsible for the conjugation of reduced glutathione to a wide number of exogenous and endogenous hydrophobic electrophiles. The amino acid sequence GST is highly conserved in most organisms including mammals. GSTs proteins are typically homodimeric, with both heterologous GST dimers have been observed. GST monomers have an average molecular weight of approximately 25-28 kDa in size. Note a different form of non-human GST (Glutathione-S-Transferase) is used as a protein expression tag commonly in molecular biology applications. All anti-GST antibodies may not react with recombinant GST-fusion proteins.

Synonyms: mouse anti-GSTK1 antibody, mouse anti-GST-K1 antibody, GST, Glutathione S-Transferases, K1 antibody, GSTK1

Note: Anti-GSTK1 antibody is tested by in ELISA and Western Blot. Suitable for most immunological techniques requiring high titer binding and lot-to-lot consistency. Specific conditions for reactivity should be optimized by the end user.

Product images:



Western Blot of Mouse anti-GSTK1 Monoclonal Antibody. Lane 1: Recombinant GSTK1 protein. Lane 2: GST. Load: 50 ng per lane. Primary antibody: Mouse anti-GSTK1 Monoclonal Antibody at 1:1,000 overnight at 4°C. Secondary antibody: HRP Mouse Secondary Antibody at 1:40,000 for 30 min at RT. Block: MB-070 for 30 min at RT. Predicted/Observed size: 27 kDa, 27 kDa for GSTK1.