

## **Product datasheet for TA396788S**

## MAP2K1 Mouse Monoclonal Antibody [Clone ID: 13B6.G12]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: 13B6.G12
Applications: ELISA, WB

Recommended Dilution: WB: 1µg/mL

**ELISA**: 1:40,000

Reactivity: Human, Mouse, Rat

**Host:** Mouse

**Isotype:** IgG1, kappa **Clonality:** Monoclonal

Immunogen: Anti-MEK1 Monoclonal Antibody was produced in mice by repeated immunizations with

synthetic peptide corresponding to amino acid residues near the C-terminus conjugated to

KLH.

**Specificity:** This protein A purified mouse monoclonal antibody reacts specifically with human MEK1.

Anti-MEK1 is purified from tissue culture supernatant by protein A purification. Cross reactivity is expected to occur with human, mouse, and rat based on sequence identity of the

peptide immunogen. This antibody does not react with the MEK2 isoform.

**Formulation:** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

**Concentration:** 1.0 mg/ml - lot specific

Conjugation: Unconjugated

Storage: Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of

reagent (25  $\mu$ L). To minimize loss of volume dilute 1:10 by adding 225  $\mu$ L of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing

and thawing.

**Stability:** Expiration date is one (1) year from date of receipt.

Database Link: Q02750



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

MEK1 antibodies detect the MEK1 isoform. Mitogen-activated protein kinase kinase 1, also known as MKK or MEK1, is an integral component of the MAP kinase cascade that regulates cell growth and differentiation. This pathway also plays a key role in synaptic plasticity in the brain. Activated MEK 1 acts as a dual specificity kinase phosphorylating both a threonine and a tyrosine residue on MAP kinase. MEK1 and MEK2 are about 80% identical to each other, and nearly identical within the kinase domain. The MEK1 antibody is ideal for investigators involved in Neuroscience, Cell Signaling and Cancer Research.


Synonyms:

mouse anti-MEK1 antibody, MAP2K, MEK, MEK 1, MKK1, PRKMK1, MEK-1 Antibody

Note:

Anti-MEK 1 (MOUSE) antibody is tested in ELISA and Western Blotting. Specific conditions of reactivity should be optimized by the end user. Expect a band of approximately 43.5 kDa.

## **Product images:**



Western Blot of Anti-MEK1 Antibody. Lane 1: MEK-1 recombinant protein. Lane 2: MEK-2 recombinant protein. Load: 50ng per lane. Primary Antibody: Anti-MEK1 supernatant clone neat over night at 4°C. Secondary Antibody: Antimouse HRP at 1:40,000 dilution.