

## Product datasheet for **TA396800S**

### **CALCA Monoclonal Antibody [Clone ID: 4C8.H6.D4]**

#### **Product data:**

Product Type:	Primary Antibodies
Clone Name:	4C8.H6.D4
Applications:	ELISA, WB
Recommended Dilution:	<b>WB:</b> 1:1,000 - 1:2,000 <b>ELISA:</b> User Optimized
Reactivity:	Human
Isotype:	IgG1, lambda
Clonality:	Monoclonal
Immunogen:	Procalcitonin Antibody was produced in mice prepared by repeated immunizations with full-length recombinant human Procalcitonin protein.
Specificity:	Anti-Procalcitonin Antibody was purified from concentrated tissue culture supernate by Protein A chromatography. This antibody is specific for human Procalcitonin. Cross-reactivity with Procalcitonin from other sources has not been determined.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Concentration:	1.0 mg/ml - lot specific
Conjugation:	Unconjugated
Storage:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Stability:	Expiration date is one (1) year from date of receipt.
Database Link:	<u><a href="#">P06881</a></u>

[View online »](#)

**Background:**


Anti-Procalcitonin antibody detects human Procalcitonin. Procalcitonin is a peptide hormone mainly produced by the C cells of the thyroid and certain endocrine cells of the lung. Under normal expression conditions, procalcitonin is immediately cleaved into three specific fragments, a N terminal residue, calcitonin and katacalcin. Levels of unprocessed procalcitonin rise significantly after bacterial infection, trauma or shock. This gene encodes the peptide hormones calcitonin, calcitonin gene-related peptide and katacalcin by tissue-specific alternative RNA splicing of the gene transcripts and cleavage of inactive precursor proteins. Calcitonin is involved in calcium regulation and acts to regulate phosphorus metabolism. Calcitonin gene-related peptide functions as a vasodilator while katacalcin is a calcium-lowering peptide. Multiple transcript variants encoding different isoforms have been found for this gene.

**Synonyms:**

mouse anti-procalcitonin antibody, calcitonin isoform CT preproprotein, calcitonin-related polypeptide alpha, CT, KC, CGRP, CALC1, CGRP1, CGRP-I

**Note:**

Anti-Procalcitonin [4C8.H6.D4] antibody was tested by ELISA and Western Blot. Specific conditions for reactivity should be optimized by the end user.

**Product images:**


Western Blot of Mouse anti-Procalcitonin Antibody. Lane 1: rProcalcitonin. Load: 50 ng. Primary antibody: Mouse anti-Procalcitonin at 1:1,000 overnight at 4°C. Secondary antibody: Peroxidase mouse secondary antibody at 1:40,000 for 30 min at RT. Block: MB-070 for 30 min at RT. Predicted/Observed size: 13.9 kDa, 13.9 kDa