

Product datasheet for **TA396862**

Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, IHC, WB
Recommended Dilution:	WB: 1:3,000 - 1:12,000 IHC: 1:1,500 - 1:5,000 ELISA: 1:200,000
Reactivity:	Streptomyces avidinii
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Streptavidin [Streptomyces avidinii]
Specificity:	Anti-Streptavidin antibody is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum as well as purified and partially purified Streptavidin [Streptomyces avidinii]. No cross reactivity occurs against Avidin.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Concentration:	1.0 mg/mL - lot specific
Conjugation:	Unconjugated
Storage:	Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.
Stability:	Expiration date is three (3) months from date of receipt.



[View online »](#)

Background:


Streptavidin antibody is a 60 kDa protein purified from the bacterium *Streptomyces avidinii*. Streptavidin homo-tetramers have an extraordinarily high affinity for biotin (also known as vitamin B7). With a dissociation constant (Kd) on the order of $\approx 10^{-14}$ mol/L,[1] the binding of biotin to streptavidin is one of the strongest non-covalent interactions known in nature. Streptavidin is used extensively in molecular biology and bio-nanotechnology due to the streptavidin-biotin complex's resistance to organic solvents, denaturants (e.g. guanidinium chloride), detergents (e.g. SDS, Triton), proteolytic enzymes, and extremes of temperature and pH. Anti-STREPTAVIDIN antibody is ideal for investigators involved in GFP and Epitope research.

Synonyms:


rabbit anti-Streptavidin Antibody, rabbit anti Streptavidin

Note:

Anti-Streptavidin antibody has been tested by ELISA and dot blot and is suitable for western blot, and immunohistochemistry, as well as other assays requiring lot-to-lot consistency.

Product images:

Rockland Rabbit anti Streptavidin (200-4195 lot 23495) and Biotin conjugated Rabbit anti-trypsin inhibitor antibody (200-4679 lot 6594) were used to detect target proteins Trypsin Inhibitor (left) and Streptavidin (right) under reducing (R) and non-reducing (NR) conditions. Reduced samples of purified target proteins contained 4% BME and were boiled for 5 minutes. Samples of ~ 1 ug of protein per lane were run by SDS-PAGE. Protein was transferred to nitrocellulose and probed with 1:1000 dilution of primary antibody (ON 4 C). Detection shown was using Dylight 649 conjugated Donkey anti rabbit (611-743-127 lot 20831 1:10K 1.5 hr RT in MB-070) and imaged on the BioRad VersaDoc System.



Rockland Rabbit anti Streptavidin (200-4195 lot 23495) and Biotin conjugated Rabbit anti-trypsin inhibitor antibody (200-4679 lot 6594) were used to detect target proteins Trypsin Inhibitor (left) and Streptavidin (right) under reducing (R) and non-reducing (NR) conditions. Reduced samples of purified target proteins contained 4% BME and were boiled for 5 minutes. Samples of ~ 1 ug of protein per lane were run by SDS-PAGE. Protein was transferred to nitrocellulose and probed with 1:1000 dilution of primary antibody (ON 4 C). Detection shown was using Dylight 649 conjugated Donkey anti rabbit (611-743-127 lot 20831 1:10K 1.5 hr RT in MB-070) and imaged on the BioRad VersaDoc System.