

Product datasheet for TA396912S

DEX Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, IP, WB
Recommended Dilution:	WB: 1:500 - 1:5,000 ELISA: 1:2,000 - 1:10,000
Reactivity:	Penicillium
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Dextranase [Penicillium species] DEX
Specificity:	Anti-Dextranase is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Rabbit Serum as well as purified and partially purified Dextranase [Penicillium species]. Cross reactivity against Dextranase from other sources is unknown.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Concentration:	1.0 mg/mL - lot specific
Conjugation:	Biotin
Storage:	Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.
Stability:	Expiration date is one (1) year from date of receipt.
Database Link:	P48845
Background:	Anti-Dextranase antibody is a transferase enzyme which catalyzes the synthesis of dextran by engaging in the endohydrolysis of 1, 6 alpha-D-glucosidic linkages in dextran. Belonging to the glycosyl hydrolase 49 family, Anti-Dextranase antibody is ideal for researchers interested in Metabolism and Signal Transduction research.




[View online »](#)

Synonyms: rabbit anti-Dextranase Antibody biotin Conjugation, biotin Conjugated rabbit anti-Dextranase Antibody, dextranase, Alpha-1,6-glucon-6-gluconohydrolase, DEX

Note: Anti-Dextranase Biotin Antibody has been tested by ELISA and western blot and is assayed against 1.0 ug of Dextranase in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:3,000 to 1:14,000 of the reconstitution concentration is suggested for this product.

Product images:



Western Blot of Anti-Dextranase Antibody Biotin Conjugated. Lane 1: Molecular Weight Marker. Lane 2: Dextranase. Primary Antibody: Rabbit Anti-Dextranase Biotin Conjugate at 1:1000 overnight at 2-8°C. Secondary Antibody: Streptavidin-HRP (p/n S000-03) at 1:40,000 for 30 min at RT. Block: BlockOut Buffer (p/n MB-073). Expected MW: ~66kDa.