

Product datasheet for **TA396930**

gox Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, IHC, WB
Recommended Dilution:	WB: 1:3500 - 1:16,000 IHC: 1:2,000 - 1:8,000 ELISA: 1:35,000 - 1:160,000
Reactivity:	Aspergillus niger
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Anti-Glucose Oxidase Antibody was produced by repeated immunizations with Aspergillus niger Glucose Oxidase protein.
Specificity:	Anti-Glucose Oxidase is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum as well as purified and partially purified Glucose Oxidase [Aspergillus niger]. Cross reactivity against Glucose Oxidase from other tissues and species may occur but have not been specifically determined.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reconstitution Method:	Restore with deionized water (or equivalent) - Reconstitution Volume: 100 µL
Concentration:	1.0 mg/ml - lot specific
Conjugation:	Unconjugated
Storage:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Stability:	Expiration date is one (1) year from date of receipt.
Database Link:	P13006




[View online »](#)

Background: Anti-Glucose Oxidase detects glucose oxidase. Glucose oxidase enzyme (GOx) is an oxidoreductase that catalyzes the oxidation of glucose to hydrogen peroxide and D-glucono- δ -lactone. In cells, it aids in breaking the sugar down into its metabolites. Glucose oxidase is widely used for the determination of free glucose in body fluids, in vegetal raw material, and in the food industry. It also has many applications in biotechnologies, typically enzyme assays for biochemistry including biosensors in nanotechnologies. Anti-Glucose Oxidase Antibody is ideal for investigators involved in Cell Signaling and Signal Transduction research.

Synonyms: rabbit anti-Glucose Oxidase Antibody, Beta D Glucose Oxygen 1 Oxido Reductase antibody, Glucose oxidase (Precursor) antibody, Glucose Oxyhydrase antibody, GOD antibody

Note: Anti-Glucose Oxidase Antibody has been tested by ELISA and western blot and is suitable for IHC. Researchers should determine optimal titers for applications that are not stated below.

Product images:



Western Blot of Rabbit anti-Glucose Oxidase Antibody. Lane 1: Glucose Oxidase. Load: 50 ng per lane. Primary antibody: Glucose Oxidase antibody at 1:1,000 for overnight at 4°C. Secondary antibody: DyLight™ 649 rabbit secondary antibody (p/n 611-143-002) at 1:20,000 for 30 min at RT. Block: (p/n MB-070) for 30 min at RT. Predicted/Observed size: 80 kDa, 80 kDa for Glucose Oxidase from *Aspergillus niger*. Other band(s): None.