

Product datasheet for TA396963S

PRSS2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, WB

Recommended Dilution: WB: 1:500 - 1:3,000

ELISA: 1:5,000 - 1:25,000

Host: Rabbit

Clonality: Polyclonal

Immunogen: Trypsinogen [Bovine Pancreas]

Specificity: Anti-Trypsinogen is an IgG fraction antibody purified from monospecific antiserum by a

multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum as well as

purified and partially purified Trypsinogen [Bovine Pancreas]. Cross reactivity against Trypsinogen from other tissues and species may occur but have not been specifically

determined.

Formulation: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Concentration: 1 mg/ml - lot specific

Conjugation: Unconjugated

Storage: Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of

reagent (25 μ L). To minimize loss of volume dilute 1:10 by adding 225 μ L of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing

and thawing.

Stability: Expiration date is one (1) year from date of receipt.

Database Link: Q29463



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



PRSS2 Rabbit Polyclonal Antibody - TA396963S

Background: Trypsinogen, is a precursor to trypsin and a digestive enzyme. It functions as storage of an

inactive form of trypsin and kept in the pancreas then released in significant amount, along with other enzymes, when required for protein digestion. Serum trypsinogen is measured using a blood test. Diseases associated with Trypsinogen high levels or mutations are acute pancreatitis and cystic fibrosis. Anti-Trypsinogen Antibody can be useful for researchers

interested in studying the serum or pancreas.

Synonyms: rabbit anti-Trypsinogen Antibody, Brain trypsinogen antibody, Cationic trypsinogen antibody,

Mesotrypsinogen antibody, Pancreatic anionic trypsinogen antibody, Protease, serine, 2

antibody, PRSS2 antibody

Note: Trypsinogen Antibody has been tested by ELIDA and dot blot and is assayed against 1.0 ug of

Trypsinogen [Bovine Pancreas] in a standard ELISA using Peroxidase conjugated Affinity

Purified anti-Rabbit IgG [H&L] (Goat) code #611-1302 and (ABTS (2,2'-azino-bis-[3-

ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:2,300 to 1:8,400 of the reconstitution concentration is

suggested for this product.