

Product datasheet for **TA396969**

SOD1 Rabbit Polyclonal Antibody

Product data:

| | |
|------------------------|--|
| Product Type: | Primary Antibodies |
| Applications: | ELISA, WB |
| Recommended Dilution: | WB: 1:500 - 1:2,000 ELISA: 1:10,000 - 1:50,000 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Immunogen: | Superoxide Dismutase [Bovine erythrocytes] |
| Specificity: | Anti-Superoxide Dismutase (SOD) is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum as well as purified and partially purified Superoxide Dismutase [Bovine erythrocytes]. Cross reactivity against Superoxide Dismutase from other tissues and species may occur but have not been specifically determined. |
| Formulation: | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Reconstitution Method: | Restore with deionized water (or equivalent) - Reconstitution Volume: 100 µL |
| Concentration: | 1.0 mg/ml - lot specific |
| Conjugation: | Unconjugated |
| Storage: | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
| Stability: | Expiration date is one (1) year from date of receipt. |
| Database Link: | P00442 |
| Background: | Superoxide Dismutase destroys radicals which are normally produced within the cells and which are toxic to biological systems. |



[View online »](#)

Synonyms: rabbit anti-Superoxide Dismutase 1 Antibody, rabbit anti-SOD1 antibody, ALS 1 antibody, ALS antibody, ALS1 antibody, Amyotrophic lateral sclerosis 1 adult antibody, Amyotrophic lateral sclerosis 1 antibody, Cu/Zn SOD antibody, Cu/Zn superoxide dismutase antibody, Homodimer antibody, SOD 1

Note: Superoxide Dismutase (SOD) Antibody has been tested by western blot and is suitable to be assayed against 1.0 ug of Superoxide Dismutase [Bovine erythrocytes] in a standard ELISA using Peroxidase conjugated Affinity Purified anti-Rabbit IgG [H&L] (Goat) code #611-1302 and (ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:10,000 to 1:50,000 is suggested for this product.