

Product datasheet for **TA396978S**

phoA Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	WB: 1:500 - 1:2,000 ELISA: 1:4,000 - 1:20,000
Reactivity:	Escherichia coli
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Alkaline Phosphatase [E.coli]
Specificity:	Alkaline phosphatase is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum as well as purified and partially purified Alkaline Phosphatase [E.coli]. No cross reactivity occurs against mammalian forms of Alkaline Phosphatase.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Concentration:	1.0 mg/mL - lot specific
Conjugation:	HRP
Storage:	Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.
Stability:	Expiration date is one (1) year from date of receipt.
Database Link:	P00634



[View online »](#)

Background:


Alkaline phosphatase (ALP, ALKP) is a hydrolase enzyme responsible for removing phosphate groups from many types of molecules, including nucleotides, proteins, and alkaloids. The process of removing the phosphate group is called dephosphorylation. All mammalian alkaline phosphatase isoenzymes except placental (PALP and SEAP) are inhibited by homoarginine, and, in similar manner, all except the intestinal and placental ones are blocked by levamisole. Heating for ~2 hours at 65°C inactivated most isoenzymes except Placental isoforms (PALP and SEAP).

Synonyms:

rabbit anti-Alkaline Phosphatase Antibody Peroxidase Conjugation, HRP conjugated rabbit anti-Alkaline Phosphatase Antibody, Akp2 antibody, ALP antibody, Nagao isozyme antibody, Placental alkaline phosphatase antibody, Placental like alkaline phosphatase antibody, PLAP 1 antibody, PLAP like antibody, Regan isozyme antibody, Kasahara isozyme antibody

Note:

Anti-Alkaline phosphatase Peroxidase Conjugated Antibody has been tested by ELISA and western blot and is assayed against 1.0 ug of Alkaline Phosphatase [E.coli] in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:3,500 to 1:15,000 of the reconstitution concentration is suggested for this product.

Product images:

Western Blot of Rabbit anti-Alkaline Phosphatase (E. Coli) Antibody Peroxidase Conjugated. Lane 1: Alkaline Phosphatase (E. Coli). Load: 50 ng per lane. Primary antibody: Rabbit anti-Alkaline Phosphatase (E. Coli) Antibody Peroxidase Conjugated at 1:1,000 overnight at 4°C. Secondary antibody: n/a. Block: MB-070 for 30 min at RT. Predicted/Observed size: 50 kDa, 50 kDa for Alkaline Phosphatase (E. Coli).