

## Product datasheet for **TA397001S**

### **GALM Rabbit Polyclonal Antibody**

#### **Product data:**

<b>Product Type:</b>	Primary Antibodies
<b>Applications:</b>	ELISA, IP, WB
<b>Recommended Dilution:</b>	<b>WB:</b> 1:500 - 1:5,000 <b>ELISA:</b> 1:5,000 - 1:20,000
<b>Reactivity:</b>	Porcine
<b>Host:</b>	Rabbit
<b>Clonality:</b>	Polyclonal
<b>Immunogen:</b>	Mutarotase [Porcine Kidney]
<b>Specificity:</b>	Mutarotase is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum as well as purified and partially purified Mutarotase [Porcine Kidney]. Cross reactivity against Mutarotase from other tissues and species may occur but have not been specifically determined.
<b>Formulation:</b>	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
<b>Concentration:</b>	1.0 mg/mL - lot specific
<b>Conjugation:</b>	HRP
<b>Storage:</b>	Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.
<b>Stability:</b>	Expiration date is one (1) year from date of receipt.
<b>Database Link:</b>	<a href="#">Q9GKX6</a>
<b>Background:</b>	Mutarotase is found especially in mammalian tissues that catalyzes the interconversion of anomeric forms of some sugars: α- and β- forms of d-glucose, l-arabinose, d-galactose, d-xylose, lactose, and maltose.




[View online »](#)

**Synonyms:** rabbit anti-Mutarotase Antibody Peroxidase Conjugation, HRP Conjugated rabbit anti-Mutarotase Antibody, Aldose 1 epimerase antibody, BLOCK25 protein antibody, Galactose mutarotase antibody, GALM antibody

**Note:** Anti-Mutarotase Peroxidase Conjugated Antibody has been tested by ELISA and western blot. This product is assayed against 1.0 ug of Mutarotase [Porcine Kidney] in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:500 to 1:1,500 of the reconstitution concentration is suggested for this product.

### Product images:



Western Blot of Rabbit Anti-Mutarotase Antibody Peroxidase Conjugated. Lane 1: Molecular Weight Marker. Lane 2: Mutarotase [0.05µg]. Primary Antibody: Anti-Mutarotase HRP at 1:1000 overnight at 2-8°C. Secondary Antibody: none. Blocking: BlockOut Buffer (p/n MB-073) for 30mins at RT. Predicted MW: ~38kDa.