

Product datasheet for **TA397066**

AVD Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, IHC, WB
Recommended Dilution:	WB: 1:500 - 1:2,500 IHC: 1:250 - 1,000 ELISA: 1:2,000 - 1:10,000
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Avidin (Hen Egg White)
Specificity:	Anti-Avidin Antibody Biotin Conjugated is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Rabbit Serum, as well as purified and partially purified Avidin [Hen Egg White].
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reconstitution Method:	Restore with deionized water (or equivalent) - Reconstitution Volume: 100 µL
Concentration:	1.0 mg/mL - lot specific
Conjugation:	Biotin
Storage:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Stability:	Expiration date is one (1) year from date of receipt.
Database Link:	P02701



[View online »](#)

Background:


Avidin is a glycoprotein with a molecular weight of approximately 62.4 kDa. Avidin is a biotin binding protein that shows high sequence homology in birds, reptiles and amphibians. Hen egg white avidin is a tetrameric protein composed of four identical subunits, each with the ability to bind biotin with high affinity and specificity ($K_d \sim 10^{15} M$). In biotechnology, the functional consequence of tetrameric biotin binding is signal amplification. Biotin-avidin bridging is a great way to increase signal strength while maintaining specificity. The sequence of avidin only shows 30% homology with streptavidin, and anti-avidin and anti-streptavidin antibodies are not immunologically cross reactive. Biotin is widely used throughout the biotechnology industry to conjugate proteins for biochemical assays. Biotin's small size typically does not affect the biological activity of protein upon biotinylation. Biotinylated proteins of interest can be enriched from a sample due to highly stable interactions.

Synonyms:

rabbit anti-Avidin Antibody biotin Conjugation, Biotin conjugated rabbit anti-Avidin antibody, Avidin BAC, Anti-Avidin biotin Antibody, Egg White

Note:

Anti-Avidin Antibody Biotin Conjugated has been tested by ELISA and western blot and is suitable for immunohistochemistry, immunomicroscopy as well as other antibody based assays using streptavidin or avidin conjugates requiring lot-to-lot consistency.

Product images:

Western Blot of Rabbit anti-Avidin Biotin Conjugated Antibody. Lane 1: Avidin. Lane 2: None. Load: 100 ng per lane. Primary antibody: Avidin antibody at 1:1000 for 1H at RT. Secondary antibody: Peroxidase streptavidin secondary antibody at 1:40,000 for 30 min at RT. Block: MB-070 for 30 min at RT. Predicted/Observed size: ~17 kDa, ~17 kDa for monomeric Avidin. Other band(s): None.