

## Product datasheet for **TA397078S**

### DAO Sheep Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, IP, WB
Recommended Dilution:	<b>WB:</b> 1:500 - 1:5,000 <b>ELISA:</b> 1:5,000 - 1:20,000
Reactivity:	Porcine
Host:	Sheep
Clonality:	Polyclonal
Immunogen:	D-Amino Acid Oxidase [Pig Kidney]
Specificity:	Anti-D-AMINO ACID OXIDASE is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Sheep Serum as well as purified and partially purified D-Amino Acid Oxidase [Pig Kidney]. Cross reactivity against D-Amino Acid Oxidase from other tissues and species may occur but have not been specifically determined.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Concentration:	1.0 mg/ml - lot specific
Conjugation:	Unconjugated
Storage:	Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.
Stability:	Expiration date is one (1) year from date of receipt.
Database Link:	<a href="#">P00371</a>



[View online »](#)

**Background:**


D-Amino Acid Oxidase regulates the level of the neuromodulator D-serine in the brain. It has high activity towards D-DOPA and contributes to dopamine synthesis. It could act as a detoxifying agent which removes D-amino acids accumulated during aging. It acts on a variety of D-amino acids, with a preference for those having small hydrophobic side chains, followed by those bearing polar, aromatic, and basic groups. D-Amino Acid Oxidase does not act on acidic amino acids.

**Synonyms:**

sheep anti-D-Amino Acid Oxidase Antibody, D-amino-acid oxidase, DAMOX, DAAO, DAO

**Note:**

D-AMINO ACID OXIDASE antibody has been tested by ELISA and western blot. This product is assayed against 1.0 ug of D-Amino Acid Oxidase [Pig Kidney] in a standard ELISA using Peroxidase conjugated Affinity Purified anti-Sheep IgG [H&L] (Goat) code #611-1302 and ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:20,000 to 1:60,000 of the reconstitution concentration is suggested for this product.

**Product images:**

Western Blot of Sheep anti-D-Amino Acid Oxidase Antibody. Lane 1: D-Amino Acid Oxidase. Load: 50 ng per lane. Primary antibody: D-Amino Acid Oxidase primary antibody at 1:1,000 overnight at 4°C. Secondary antibody: Peroxidase sheep secondary antibody (p/n 613-4302) at 1:40,000 for 30 min at RT. Block: (p/n MB-070) for 30 min at RT. Predicted/Observed size: 39 kDa, 39 kDa for D-Amino Acid Oxidase.