

Product datasheet for **TA397169**


Goat IgG F(c) Antibody Peroxidase Conjugated

Product data:

Product Type:	Secondary Antibodies
Product Name:	Goat IgG F(c) Antibody Peroxidase Conjugated
Applications:	ELISA, IHC, WB
Recommended Dilution:	WB: 1:1,000 - 1:5,000 IHC: 1500 - 1:2,500 ELISA: 1:5,000 - 1:25,000
Host:	Donkey
Immunogen:	Goat IgG F(c) fragment
Formulation:	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reconstitution Method:	Restore with deionized water (or equivalent) - Reconstitution Volume: 2.0 mL
Concentration:	10.0 mg/mL - lot specific
Conjugation:	HRP
Storage:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Note:	Anti-Goat IgG F(c) Peroxidase Conjugated Antibody has been tested by ELISA. This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.



[View online »](#)

Product images:

ELISA Results of Donkey Anti-Goat IgG Fc Antibody Peroxidase Conjugated. Each well was coated in duplicate with 1.0 μg of Goat IgG Fc (p/n 005-0103). The working dilution is 32,600. The starting dilution of antibody was 5 $\mu\text{g}/\text{ml}$ and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using TMB substrate (p/n TMBE-1000).