

## **Product datasheet for TA397267**

## Mouse IgM (H&L) Antibody Peroxidase Conjugated

**Product data:** 

**Product Type:** Secondary Antibodies

**Product Name:** Mouse IgM (H&L) Antibody Peroxidase Conjugated

**Applications:** ELISA, IHC, WB

Recommended Dilution: WB: 1:1,000 - 1:10,000

IHC: 1:500 - 1:2,500

**ELISA**: 1:10,000 - 1:50,000

Reactivity: Mouse Host: Rabbit

Immunogen: Anti-Mouse IgM (H&L) was produced by repeated immunization with mouse IgM whole

molecule in rabbit.

**Formulation:** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Reconstitution Method: Restore with deionized water (or equivalent) - Reconstitution Volume: 2.0 mL

**Concentration:** 10 mg/mL - lot specific

Conjugation: HRP

Storage: Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -

20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as

an undiluted liquid. Dilute only prior to immediate use.

**Note:** This secondary peroxidase conjugated antibody anti-Mouse has been tested by dot blot and

ELISA and is ideal for investigators who routinely perform titration assays, western-blot, immunoprecipitation and more generally immunoassays. Mouse IgM (H&L) peroxidase conjugated antibody has been assayed against 1.0 ug of mouse IgM in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:10,000 to 1:40,000 of

the reconstitution concentration is suggested for this product.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



Dot Blot results of Rabbit Anti-Mouse IgM Antibody Peroxidase Conjugated. Dots are Mouse IgM at (1) 100ng, (2) 33.3ng, (3) 11.1ng, (4) 3.70ng, (5) 1.23ng. Blocking: MB-070 for 30 min at RT. Primary Antibody: none. Secondary Antibody: Rabbit anti-Mouse IgM-HRP at 1µg/mL for 1hr at RT. Imaged with BioRad ChemiDoc, Chemi filter.