

Product datasheet for **TA397489**

H3C14 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ChIP, IF, IHC, WB
Recommended Dilution:	WB: 1:500 IHC: 1:100 IF: 1:100 ChIP: 2-5µg/million cells
Reactivity:	C. elegans, Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Histone H3 [Asym-dimethyl Arg17] affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic asymmetric dimethylated peptide surrounding Arginine 17 of human Histone H3.2.
Specificity:	Anti-Histone H3 [Asym-dimethyl Arg17] was affinity purified from monospecific antiserum by immunoaffinity chromatography. This antibody reacts with human Histone H3.2. A BLAST analysis was used to suggest cross-reactivity with Human, mouse, and C. elegans. Predicted to react with many species including rat, chicken, Xenopus, Drosophila, and plant based on 100% sequence homology. Cross-reactivity with Histone H3 from other sources has not been determined.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Concentration:	0.83 mg/ml - lot specific
Conjugation:	Unconjugated
Storage:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Stability:	Expiration date is one (1) year from date of receipt.
Gene Name:	histone cluster 2, H3c
Database Link:	Entrez Gene 333932 Human Entrez Gene 126961 Human Q71DI3



[View online »](#)

Background:

Chromatin is the arrangement of DNA and proteins in which chromosomes are formed. Correspondingly, chromatin is formed from nucleosomes, which are comprised of a set of four histone proteins (H2A, H2B, H3, H4) wrapped with DNA. Chromatin is a very dynamic structure in which numerous post-translational modifications work together to activate or repress the availability of DNA to be copied, transcribed, or repaired. These marks decide which DNA will be open and commonly active (euchromatin) or tightly wound to prevent access and activation (heterochromatin). Common histone modifications include methylation of lysine and arginine, acetylation of lysine, phosphorylation of threonine and serine, and sumoylation, biotinylation, and ubiquitylation of lysine. In particular, dimethylation of H3 Arg17 (H3 R17Me₂) has been linked to gene activation. Coactivator-associated arginine methyltransferase-1 (CARM1) methylates Arg17 with its protein arginine methyltransferase (PRMT) catalytic core. Activation of this modification is linked to transcription hormone response promoters, as well as cell fate regulation. Interestingly, H3 methylation of R17 and R26 contributes to greater pluripotency potential of stem cells, while downregulation of this PTM increases differentiation. Anti-Histone H3 are ideal for researchers interested in Chromatin Modifiers, Chromatin Research, Histones and Modified Histones, and Epigenetics research.

Synonyms:


rabbit anti-Histone H3 Asym-dimethyl Arg17 antibody, H3.3B, H3 histone, family 3A, H3.3AH3F3H3F3B, histone H3.3, MGC87783, MGC87782, H3R17me_{2a}

Note:

Anti-Histone H3 [Asym-dimethyl Arg17] antibody is tested for Western Blot, Chromatin Immunoprecipitation, and Immunofluorescence. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately ~15.4 kDa corresponding to Histone H3 protein by Western Blotting in HeLa histone prep lysate or the appropriate cell lysate or extract. Epi-Plus™ antibody production in collaboration with Novus Biologicals.

Protein Pathways:

Systemic lupus erythematosus

Product images:


Immunofluorescence of Rabbit Anti-Histone H3 [Asym-dimethyl Arg17] Antibody. Tissue: HeLa cells. Fixation: 0.5% PFA. Antigen retrieval: Not required. Primary antibody: Histone H3 [Asym-dimethyl Arg17] antibody at a 1:50 dilution for 1 h at RT. Secondary antibody: FITC secondary antibody at 1:10,000 for 45 min at RT. Localization: Histone H3 [Asym-dimethyl Arg17] is nuclear and chromosomal. Staining: Histone H3 [Asym-dimethyl Arg17] is expressed in green, nuclei are counterstained with Dapi (blue).