

Product datasheet for TA397508

Arhgap22 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, IF, IHC, WB

Recommended Dilution: WB: 1 µg/ml

IHC: 1:100-1:500 **IF**: 1:100-1:500

ELISA: 1:20,000 - 1:60,000

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Immunogen: ARHGAP22 affinity purified antibody was prepared from whole rabbit serum produced by

repeated immunizations with a synthetic phospho-peptide corresponding to the region

surrounding mouse pS22 region of ARHGAP22.

Specificity: Anti-ARHGAP22 pS22 was affinity purified from monospecific antiserum by immunoaffinity

chromatography. This antibody is specific for phosphorylated ARHGAP22 at Serine 22. It also recognizes the S397->A mutation but not the S22->A mutation. A BLAST analysis was used to suggest cross-reactivity with Mouse, Rat and Human based on 100% sequence homology. Cross-reactivity with ARHGAP22 pS22 from other sources has not been determined.

Formulation: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Concentration: 1.07 mg/ml - lot specific

Conjugation: Unconjugated

Storage: Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for

extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as

an undiluted liquid. Dilute only prior to immediate use.

Stability: Expiration date is six (6) months from date of receipt.

Gene Name: Rho GTPase activating protein 22

Database Link: Entrez Gene 239027 Mouse

Q8BL80



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background: ARHGAP22 is a Rho GTPase-activating protein involved in the signal transduction pathway

that regulates endothelial cell capillary tube formation during angiogenesis. It acts as a GTPase activator for RAC1 by converting it to an inactive GDP-bound state and also inhibits RAC1-dependent lamellipodia formation. It may also play a role in transcription regulation via its interaction with VEZF1, by regulating activity of the endothelin-1 (EDN1) promoter. Anti-ARHGAP22 [p Ser22] antibody is ideal for researchers interested in Diabetes Research, Lipid

and Metabolism research.


Synonyms: rabbit anti-ARHGAP22 pS22 Antibody, Rho-type GTPase-activating protein 22, RHOGAP2, Rho

GTPase activating protein 22, rho GTPase-activating protein 22, ARHGAP 22, ARHGAP-22

Note: Anti-ARHGAP22 pS22 antibody is tested in Western Blot and ELISA useful for Immunostaining.

Specific conditions for reactivity should be optimized by the end user. Expect a band approximately ~77.8 kDa corresponding to the appropriate cell lysate or extract.

Product images:



Western Blot of Rabbit anti-ARHGAP22 pS22 antibody. Lane 1: NIH3T3 cells transfected with a null vector. Lane 2: NIH3T3 cells transfected with ARHGAP22. Lane 3: NIH3T3 cells transfected with ARHGAP22 S22 to alanine mutation. Lane 4: NIH3T3 cells transfected with ARHGAP22 S397 to alanine mutaion. Primary antibody: Left: ARHGAP22 pS22, Right: ARHGAP22 pS397 antibody at 1µg/mL for overnight at 4°C. Secondary antibody: IRDye800™ rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO O/N at 4°C. Predicted/Observed size: 68 kDa for ARHGAP22. Other band(s): Unmodified ARHGAP22. ARHGAP22-pS22 antibody recognizes the S397>A mutation, not the S22>mutation; ARHGAP22 pS397 recognizes the pS22>A mutation, not the pS397>A mutation; Confirms the specificity of each ARHGAP22 phospho specific antibody.