

Product datasheet for **TA397529**

AR Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	WB: 1ug/ml ELISA: 1: 10,000
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Anti-Androgen Receptor pY267 affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to the internal region of human ANDR protein.
Specificity:	Anti-Androgen Receptor pY267 was affinity purified from monospecific antiserum by immunoaffinity chromatography. A BLAST analysis was used to suggest cross-reactivity with human and chimpanzee based on 100% sequence homology. Cross-reactivity with Androgen Receptor pY267 from other sources has not been determined.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Concentration:	1.1 mg/mL - lot specific
Conjugation:	Unconjugated
Storage:	Store AR Antibody at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Stability:	Expiration date is one (1) year from date of receipt.
Database Link:	P10275



[View online »](#)

- Background:** Androgen Receptor detects human Androgen receptor. The androgen receptor (AR) is an ~110 kDa androgen-dependent transcription factor that is a member of the steroid/nuclear receptor gene superfamily. The AR signaling pathway plays a key role in development and function of male reproductive organs, including the prostate and epididymis. AR also plays a role in nonreproductive organs, such as muscle, hair follicles, and brain. Abnormalities in the AR signaling pathway have been linked to a number of diseases, including prostate cancer, Kennedy's disease and male infertility. The PI3K/Akt signaling pathway plays an important role in regulating AR activity through phosphorylation of AR at Ser213/210 and Ser791/790. Growth factors or cytokines may induce phosphorylation of AR through the PI3K/Akt pathway. Activation of the PI3K/Akt pathway is thought to have a survival role in prostate cancer by protecting cells from apoptosis. Anti-Androgen Receptor Antibody is ideal for investigators involved in neuroscience, cytokines and growth factors, and transcription factors.
- Synonyms:** rabbit Anti-Androgen receptor pY267, Dihydrotestosterone receptor, Nuclear receptor subfamily 3 group C member 4, AR, DHTR, NR3C4
- Note:** Anti-Androgen Receptor antibody is useful for ELISA, Immunohistochemistry, and Western Blot. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately ~99kDa corresponding to the appropriate cell lysate or extract.