

Product datasheet for **TA397622**


Dog IgG F(c) Antibody Biotin Conjugated

Product data:

Product Type:	Secondary Antibodies
Product Name:	Dog IgG F(c) Antibody Biotin Conjugated
Applications:	ELISA, IHC, WB
Recommended Dilution:	WB: 1:2,000 - 1:10,000 IHC: 1:1,000 - 1:5,000 ELISA: 1:20,000 - 1:100,000
Host:	Goat
Immunogen:	Dog IgG F(c) fragment
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reconstitution Method:	Restore with deionized water (or equivalent) - Reconstitution Volume: 1.0 mL
Concentration:	2.0 mg/mL - lot specific
Conjugation:	Biotin
Storage:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Note:	Anti-DOG IgG F(c) Biotin Conjugated Antibody has been tested by ELISA and is assayed against 1.0 ug of Dog IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:40,000 to 1:130,000 is suggested for this product.



[View online »](#)

Product images:

ELISA Results of Goat Anti-Dog IgG F(c) Antibody Biotin Conjugated tested against purified Dog IgG F(c). Each well was coated in duplicate with 1.0 μg of Dog IgG F(c) (p/n 004-0103). The working dilution is 1:88,000. The starting dilution of antibody was 5 $\mu\text{g}/\text{ml}$ and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using HRP stabilizer (p/n MB-076), Streptavidin HRP conjugated (p/n S000-03) and TMB substrate (p/n TMBE-1000).