

Product datasheet for **TA397642**

Goat IgG (H&L) Antibody


Product data:

Product Type:	Secondary Antibodies
Product Name:	Goat IgG (H&L) Antibody
Applications:	ELISA, IHC, WB
Recommended Dilution:	WB: 1:2,000 - 1:10,000 IHC: 1:1,000 - 1:5,000 ELISA: 1:23,400-1:43,400
Reactivity:	Goat
Host:	Donkey
Immunogen:	Anti-Goat IgG (H&L) was produced by repeated immunization with goat IgG whole molecule in donkey
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Concentration:	2.0 mg/mL - lot specific
Conjugation:	Unconjugated
Storage:	Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.
Note:	Antibody Anti-Goat IgG (H&L) has been tested by ELISA and western blot. This antibody is suitable for immunoblotting (western or dot blot), ELISA, immunoelectron microscopy and immunohistochemistry as well as other antibody-based enzymatic assays requiring lot-to-lot consistency.



[View online »](#)

Product images:



Western Blot of Donkey anti-Goat IgG antibody.
Lane 1: Goat IgG (p/n 005-0102). Lane 2: none.
Load: 50 ng per lane. Primary antibody: Goat IgG antibody at 1:1000 for overnight at 4°C.
Secondary antibody: Peroxidase donkey secondary antibody at 1:40,000 for 30 min at RT.
Block: MB-070 for 30 min at RT.
Predicted/Observed size: 28 and 55 kDa for Goat IgG. Other band(s): none.