

## Product datasheet for **TA397644**


### Goat IgG (H&L) Antibody Peroxidase Conjugated

#### Product data:

Product Type:	Secondary Antibodies
Product Name:	Goat IgG (H&L) Antibody Peroxidase Conjugated
Applications:	ELISA, IHC, WB
Recommended Dilution:	<b>WB:</b> 1:1,000 - 1:5,000 <b>IHC:</b> 1:500 - 1:2,500 <b>ELISA:</b> 1:10,000 - 1:150,000
Reactivity:	Goat
Host:	Donkey
Immunogen:	Anti-Goat IgG (H&L) was produced by repeated immunization with Goat IgG whole molecule in donkey.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reconstitution Method:	Restore with deionized water (or equivalent) - Reconstitution Volume: 1.0 mL
Concentration:	2.0 mg/mL - lot specific
Conjugation:	HRP
Storage:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Note:	Anti-Goat IgG (H&L) is suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. It is also suitable in IHC. Specific conditions for reactivity and signal detection should be optimized by the end user.



[View online »](#)

**Product images:**

ELISA results of purified Donkey anti-Goat IgG antibody Peroxidase conjugated tested against purified Goat IgG. Each well was coated in duplicate with 1.0 µg of Goat IgG (p/n 005-0102-0010). The starting dilution of antibody was 5µg/ml and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using 3% fish gelatin as blocking buffer, and TMB substrate (p/n TMBE-1000).