

Product datasheet for TA397666

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

Guinea Pig IgG F(c) Antibody Peroxidase Conjugated

Product data:

Product Type: Secondary Antibodies

Product Name: Guinea Pig IgG F(c) Antibody Peroxidase Conjugated

Applications: ELISA, IHC, WB

Recommended Dilution: WB: 1:1,000 - 1:10,000

IHC: 1:500 - 1:2,500

ELISA: 1:10,000 - 1:50,000

Host: Goa

Immunogen: Guinea Pig IgG F(c) fragment

Formulation: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Reconstitution Method: Restore with deionized water (or equivalent) - Reconstitution Volume: 1.0 mL

Concentration: 2.0 mg/mL - lot specific

Conjugation: HRP

Storage: Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -

20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as

an undiluted liquid. Dilute only prior to immediate use.

Note: Anti-Guinea Pig IgG F(c) Peroxidase Conjugated Antibody has been tested by ELISA. This

product has been assayed against 1.0 ug of Guinea Pig lgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for


30 minutes at room temperature. A working dilution of 1:4,000 to 1:20,000 of the

reconstitution concentration is suggested for this product.





Product images:



ELISA Results of Purified Goat Anti-Guinea Pig IgG F(c) Antibody Peroxidase Conjugated tested against purified Guinea Pig IgG F(c). Each well was coated in duplicate with 10 µg of Guinea Pig IgG F(c) (p/n 006-0103). The working dilution of Guinea Pig IgG F(c) HRP is 1:31,000. The starting dilution of antibody was 5µg/ml and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using 3% Fish Gel/PBS Blocking buffer (p/n MB-066), Guinea Pig IgG F(c) HRP conjugated and TMB substrate (p/n TMBE-1000).