

Product datasheet for **TA397805S**

Mouse IgG (H&L) Antibody Rhodamine Conjugated


Product data:

Product Type:	Secondary Antibodies
Product Name:	Mouse IgG (H&L) Antibody Rhodamine Conjugated
Applications:	FC, IF
Recommended Dilution:	IF: 1:1,000 - 1:5,000 FC: 1:500 - 1:2,500 FLISA: 1:10,000 - 1:50,000
Reactivity:	Mouse
Host:	Goat
Immunogen:	Mouse IgG whole molecule
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reconstitution Method:	Restore with deionized water (or equivalent) - Reconstitution Volume: 500 µL
Concentration:	1.0 mg/mL - lot specific
Conjugation:	TRITC
Storage:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Note:	Anti-Mouse IgG Rhodamine Antibody has been tested by dot blot and western blot and is available in a variety of formats. Anti IgG secondary antibody conjugated is suitable for ELISA, Immunohistochemistry western blotting as well as other peroxidase antibody based assays.




[View online »](#)

Product images:



Western Blot of Rhodamine Conjugated Goat anti-Mouse IgG Secondary Antibody. Lane 1: Mouse IgG. Lane 2: none. Load: 50 ng per lane. Primary antibody: none. Secondary antibody: Rhodamine goat secondary antibody at 1:1,000 for 60 min at RT. Block: MB-070 for 30 min at RT. Predicted/Observed size: 25 & 55 kDa, 25 & 55 kDa for Goat IgG. Other band(s): none.



Western Blot of Rhodamine Conjugated Goat anti-Mouse IgG Secondary Antibody. Lane 1: Mouse IgG. Lane 2: none. Load: 50 ng per lane. Primary antibody: none. Secondary antibody: Rhodamine goat secondary antibody at 1:1,000 for 60 min at RT. Block: MB-070 for 30 min at RT. Predicted/Observed size: 25 & 55 kDa, 25 & 55 kDa for Goat IgG. Other band(s): none.