

## Product datasheet for **TA397937**


### Rabbit IgG (H&L) Antibody ATTO 488 Conjugated Pre-Adsorbed

#### Product data:


|                        |  |
|------------------------|--|
| Product Type:          | Secondary Antibodies   |
| Product Name:          | Rabbit IgG (H&L) Antibody ATTO 488 Conjugated Pre-Adsorbed   |
| Applications:          | IF, WB   |
| Recommended Dilution:  | <b>WB:</b> >1:10,000<br><b>IF:</b> >1:5,000<br><b>FLISA:</b> >1:20,000   |
| Reactivity:            | Rabbit   |
| Host:                  | Goat   |
| Immunogen:             | Rabbit IgG whole molecule  |
| Formulation:           | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2   |
| Reconstitution Method: | Restore with deionized water (or equivalent) - Reconstitution Volume: 500 µL   |
| Concentration:         | 1.0 mg/mL - lot specific   |
| Conjugation:           | ATTO 488   |
| Storage:               | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.  |
| Note:                  | Anti-Rabbit IgG (H&L) conjugated to ATTO 488 has been tested by dot blot and western blot and is designed for STED microscopy, FRET, immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. The emission spectra for this ATTO conjugate matches the principle output wavelengths of most common fluorescence instrumentation. |



[View online »](#)

**Product images:**

ATTO® dyes can be used for multicolor immunofluorescent detection with low background and high signal. Examples shown are: A. Tubulin in PtK2- male Rat Kangaroo Kidney Epithelial Cells was detected using ATTO 532 labeled secondary antibody. B. Muscle alpha-actin was stained with a mouse primary antibody and ATTO 488 anti-mouse IgG (green) while Cytokeratin was stained with polyclonal rabbit anti-cytokeratin and ATTO 647N anti-rabbit IgG (red). C. HUVEC (Human umbilical vein endothelial cells) were stained with anti- Vimentin-ATTO 532 (green), anti-E-Cadherin-ATTO 655 (red) and DAPI (blue). D. Rat colon sections were stained with Anti-Aquaporin 3-ATTO 594 antibody. Hoechst 33342 (blue) is used as counterstain. Images provided courtesy of Dr. Jörg Reichwein, ATTO-TEC GmbH



ATTO® dyes can be used for multicolor immunofluorescent detection with low background and high signal. Examples shown are: A. Tubulin in PtK2- male Rat Kangaroo Kidney Epithelial Cells was detected using ATTO 532 labeled secondary antibody. B. Muscle alpha-actin was stained with a mouse primary antibody and ATTO 488 anti-mouse IgG (green) while Cytokeratin was stained with polyclonal rabbit anti-cytokeratin and ATTO 647N anti-rabbit IgG (red). C. HUVEC (Human umbilical vein endothelial cells) were stained with anti- Vimentin-ATTO 532 (green), anti-E-Cadherin-ATTO 655 (red) and DAPI (blue). D. Rat colon sections were stained with Anti-Aquaporin 3-ATTO 594 antibody. Hoechst 33342 (blue) is used as counterstain. Images provided courtesy of Dr. Jörg Reichwein, ATTO-TEC GmbH