

Product datasheet for TA397997

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Rat IgG (H&L) Antibody Rhodamine Conjugated Pre-Adsorbed

Product data:

Product Type: Secondary Antibodies

Product Name: Rat IgG (H&L) Antibody Rhodamine Conjugated Pre-Adsorbed

Applications: FC, IF

Recommended Dilution: IF: 1:1,000 - 1:5,000

FC: 1:500 - 1:2,500

FLISA: 1:10,000 - 1:50,000

Host: Goa

Immunogen: Rat IgG whole molecule

Formulation: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Reconstitution Method: Restore with deionized water (or equivalent) - Reconstitution Volume: 1.0 mL

Concentration: 2.0 mg/mL - lot specific

Conjugation: TRITC

Storage: Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -

20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as

an undiluted liquid. Dilute only prior to immediate use.


Note: This product is designed for immunofluorescence microscopy, fluorescence based plate

assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex

analysis, including multicolor imaging, utilizing various commercial platforms.



Product images:



Western Blot of Anti-Rat IgG (H&L) (GOAT) Antibody (p/n 612-1102). Lane M: 3 µl Molecular Ladder. Lane 1: Rat IgG whole molecule (p/n 012-0102). Lane 2: Rat IgG F(c) Fragment (p/n 012-0103). Lane 3: Rat IgG Fab Fragment (p/n 012-0105). Lane 4: Rat IgM Whole Molecule (p/n 012-0107). Lane 5: Rat Serum (p/n [D310-05]). All samples were reduced. Load: 50 ng per lane. Block: MB-070 for 30 min at RT. Primary Antibody: Anti-Rat IgG (H&L) (GOAT) Antibody (p/n 612-1102) 1:1,000 for 60 min at RT. Secondary Antibody: Anti-Goat IgG (DONKEY) Peroxidase Conjugated Antibody (p/n CUST10) 1:40,000 in MB-070 for 30 min at RT. Predicted/Obsevered Size: 25 and 55 kDa for Rat IgG and Serum, 25 kDa for F(c) and Fab, 78 and 25 kDa for IgM. Rat F(c) migrates slightly higher.