

Product datasheet for TA398000

Rat IgG F(c) Antibody

Product data:

Product Type: Secondary Antibodies
Product Name: Rat IgG F(c) Antibody

Applications: ELISA, IHC, WB

Recommended Dilution: WB: 1:2,000 - 1:10,000

IHC: 1:1,000 - 1:5,000 **ELISA**: 1:8,000 - 1:9,000

Host: Goat

Immunogen: Rat IgG F(c) fragment

Formulation: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Concentration: 2.14 mg/mL - lot specific

Conjugation: Unconjugated

Storage: Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an

undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents

and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Note: Anti-Rat IgG F(c) antibody has been tested by ELISA, dot blot, and western blot and is suitable

for immunohistochemistry, as well as other assays requiring lot-to-lot consistency.


OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



Western blot of Goat Anti-Rat IgG F(c) secondary antibody. Lane 1: Rat Fc (p/n 012-0103). Lane 2: None. Load: 50 ng per lane. Primary antibody: Rat IgG F(c) antibody at 1:1,000 for overnight at &C. Secondary antibody: Peroxidase goat secondary antibody at 1:40,000 for 30 min at RT. Blocking: MB-070 for 30 min at RT. Predicted/Observed size: 28 kDa, 28 kDa for Rat IgG F(c). Other band(s): None.