

## **Product datasheet for TA398009**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Rat IgG (H&L) Antibody Peroxidase Conjugated Pre-Adsorbed

**Product data:** 

**Product Type:** Secondary Antibodies

**Product Name:** Rat IgG (H&L) Antibody Peroxidase Conjugated Pre-Adsorbed

**Applications:** ELISA, IHC, WB

**Recommended Dilution: WB**: 1:5,000 - 1:25,000

**IHC**: 1:1,000 - 1:5,000

**ELISA**: 1:50,000 - 1:350,000

Host: Goat

Immunogen: Rat IgG whole molecule

**Formulation:** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Reconstitution Method: Restore with deionized water (or equivalent) - Reconstitution Volume: 1.0 mL

**Concentration:** 2.0 mg/mL - lot specific

Conjugation: HRP

Storage: Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -

20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as


an undiluted liquid. Dilute only prior to immediate use.

**Note:** Anti-Rat IgG antibody is suitable for use in ELISA, immunohistochemistry, and western blot.

Specific conditions for reactivity should be optimized by the end user.



## **Product images:**



Western Blot of Anti-Rat IgG (H&L) (GOAT) Antibody (Min X Human Serum Proteins) (p/n 612-1125). Lane M: 3 µl Molecular Ladder. Lane 1: Rat IgG whole molecule (p/n 012-0102). Lane 2: Rat IgG F(c) Fragment (p/n 012-0103). Lane 3: Rat IgG Fab Fragment (p/n 012-0105). Lane 4: Rat IgM Whole Molecule (p/n 012-0107). Lane 5: Rat Serum (p/n [D310-05]). All samples were reduced. Load: 50 ng per lane. Block: MB-070 for 30 min at RT. Primary Antibody: Anti-Rat IgG (H&L) (GOAT) Antibody (Min X Human Serum Proteins) (p/n 612-1125) 1:1,000 for 60 min at RT. Secondary Antibody: Anti-Goat IgG (DONKEY) Peroxidase Conjugated Antibody (p/n CUST10) 1:40,000 in MB-070 for 30 min at RT. Predicted/Obsevered Size: 25 and 55 kDa for Rat IgG and Serum, 25 kDa for F(c) and Fab, 78 and 25 kDa for IgM. Rat F(c) migrates slightly higher.