

Product datasheet for TA398039

Rat IgG (H&L) Antibody Pre-adsorbed

Product data:

Product Type: Secondary Antibodies

Product Name: Rat IgG (H&L) Antibody Pre-adsorbed

Applications: ELISA, IHC, WB

Recommended Dilution: WB: 1:2,000 - 1:10,000

IHC: 1:1,000 - 1:5,000

ELISA: 1:60,000 - 1:300,000

Host: Donkey

Immunogen: Rat IgG whole molecule

Formulation: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Concentration: 1.12 mg/mL - lot specific

Conjugation: Unconjugated

Storage: Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an

undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents

and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Note: Anti-Rat IgG antibody has been tested in western blot and is suitable for use in ELISA,

immunohistochemistry, and western blot. Specific conditions for reactivity should be

optimized by the end user.




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



Western Blot of Anti-Rat IgG (H&L) (DONKEY) Antibody (Min X Bv Ch Gt GP Ham Hs Hu Ms Rb & Sh Serum Proteins) (p/n 612-701-120). Lane M: 3 μl Molecular Ladder. Lane 1: Rat IgG whole molecule (p/n 012-0102). Lane 2: Rat IgG F(c) Fragment (p/n 012-0103). Lane 3: Rat IgG Fab Fragment (p/n 012-0105). Lane 4: Rat IgM Whole Molecule (p/n 012-0107). Lane 5: Rat Serum (p/n [D310-05]). All samples were reduced. Load: 50 ng per lane. Block: MB-070 for 30 min at RT. Primary Antibody: Anti-Rat IgG (H&L) (DONKEY) Antibody (Min X Bv Ch Gt GP Ham Hs Hu Ms Rb & Sh Serum Proteins) (p/n 612-701-120) 1:10,000 for 60 min at RT. Secondary Antibody: Anti-Donkey IgG (GOAT) Peroxidase Conjugated Antibody (p/n 616-1302) 1:40,000 in MB-070 for 30 min at RT. Predicted/Obsevered Size: 25 and 55 kDa for Rat IgG and Serum, 25 kDa for F(c) and Fab, 78 and 25 kDa for IgM. Rat F(c) migrates slightly higher.