

## Product datasheet for **TA398126**

### **F(ab')<sub>2</sub> Goat IgG (H&L) Antibody Rhodamine Conjugated**

#### **Product data:**

<b>Product Type:</b>	Secondary Antibodies
<b>Product Name:</b>	F(ab') <sub>2</sub> Goat IgG (H&L) Antibody Rhodamine Conjugated
<b>Applications:</b>	FC, IF
<b>Recommended Dilution:</b>	<b>IF:</b> 1:1,000 - 1:5,000 <b>FC:</b> 1:500 - 1:2,500 <b>FLISA:</b> 1:10,000 - 1:50,000
<b>Host:</b>	Rabbit
<b>Immunogen:</b>	Goat IgG whole molecule
<b>Formulation:</b>	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
<b>Reconstitution Method:</b>	Restore with deionized water (or equivalent) - Reconstitution Volume: 500 µL
<b>Concentration:</b>	1.0 mg/mL - lot specific
<b>Conjugation:</b>	TRITC
<b>Storage:</b>	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
<b>Note:</b>	F(ab') <sub>2</sub> Anti-Goat IgG Rhodamine Antibody has been tested by western blot and is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.



[View online »](#)