

## Product datasheet for **TA398149**

### **F(ab')<sub>2</sub> Golden Syrian Hamster IgG (H&L) Antibody Phycoerythrin Conjugated**

#### **Product data:**

<b>Product Type:</b>	Secondary Antibodies
<b>Product Name:</b>	F(ab') <sub>2</sub> Golden Syrian Hamster IgG (H&L) Antibody Phycoerythrin Conjugated
<b>Applications:</b>	FC, IF
<b>Recommended Dilution:</b>	<b>IF:</b> 1:100 - 1:250 <b>FC:</b> 1:100 - 1:250
<b>Reactivity:</b>	Hamster
<b>Host:</b>	Rabbit
<b>Immunogen:</b>	Anti-Golden Syrian Hamster IgG (H&L) was produced by repeated immunization with Golden Syrian Hamster IgG whole molecule in rabbit.
<b>Formulation:</b>	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
<b>Reconstitution Method:</b>	Restore with deionized water (or equivalent) - Reconstitution Volume: 1.0 mL
<b>Concentration:</b>	0.5 mg/mL - lot specific
<b>Conjugation:</b>	Phycoerythrin
<b>Storage:</b>	Store vial at 4° C prior to restoration. Restore with deionized water (or equivalent). This product is stable at 4° C as an undiluted liquid. Dilute only prior to immediate use. Centrifuge product if not completely clear after standing at room temperature. Do not freeze after reconstitution. Store reagent in the dark. Use subdued lighting during handling and incubation of cells prior to analysis.
<b>Note:</b>	Golden Syrian Hamster IgG (H&L) Antibody is suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10 <sup>6</sup> cells in flow cytometry is approximately 1.0 µg of antibody conjugate. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.



[View online »](#)