

## Product datasheet for **TA398187**

### F(ab')<sub>2</sub> Rabbit IgG (H&L) Antibody

#### Product data:

Product Type:	Secondary Antibodies
Product Name:	F(ab') <sub>2</sub> Rabbit IgG (H&L) Antibody
Applications:	ELISA, IHC, WB
Recommended Dilution:	<b>WB:</b> 1:2,000-1:10,000 <b>IHC:</b> 1:1,000-1:5,000 <b>ELISA:</b> 1:20,000 - 1:100,000
Reactivity:	Rabbit
Host:	Goat
Immunogen:	Anti-Rabbit IgG was produced by repeated immunization with Rabbit IgG whole molecule in goat.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Concentration:	1.0 mg/mL - lot specific
Conjugation:	Unconjugated
Storage:	Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.
Note:	F(ab') <sub>2</sub> Rabbit IgG (H&L) Antibody has been tested by ELISA and is suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10 <sup>6</sup> cells in flow cytometry is approximately 1.0 µg of antibody. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.



[View online »](#)