

## Product datasheet for **TA398263**

### **F(Ab')<sub>2</sub> Fragment Of Hamster IgG (H&L) Antibody Rhodamine Conjugated Pre-Adsorbed**

#### **Product data:**

|                               |   |
|-------------------------------|---|
| <b>Product Type:</b>          | Secondary Antibodies  |
| <b>Product Name:</b>          | F(Ab') <sub>2</sub> Fragment Of Hamster IgG (H&L) Antibody Rhodamine Conjugated Pre-Adsorbed  |
| <b>Applications:</b>          | FC, IF, IHC   |
| <b>Recommended Dilution:</b>  | <b>IHC:</b> 1:1000-1:5000<br><b>IF:</b> 1:500-1:2,500<br><b>FC:</b> 1:500-1:2,500<br><b>FLISA:</b> 1:10,000-1:50,000  |
| <b>Host:</b>                  | Goat  |
| <b>Immunogen:</b>             | Golden Syrian and Armenian Hamster IgG whole molecules  |
| <b>Formulation:</b>           | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |
| <b>Reconstitution Method:</b> | Restore with deionized water (or equivalent) - Reconstitution Volume: 500 µL  |
| <b>Concentration:</b>         | 1.0 mg/mL - lot specific  |
| <b>Conjugation:</b>           | TRITC   |
| <b>Storage:</b>               | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
| <b>Note:</b>                  | F(ab') <sub>2</sub> Anti-Golden Syrian & Armenian Hamster IgG Rhodamine Antibody has been tested by dot blot and is suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency.  |



[View online »](#)