

## Product datasheet for **TA398265**

### **F(Ab')<sub>2</sub> Fragment Of Hamster IgG (H&L) Antibody Fluorescein Conjugated Pre-Adsorbed**


#### **Product data:**

<b>Product Type:</b>	Secondary Antibodies
<b>Product Name:</b>	F(Ab') <sub>2</sub> Fragment Of Hamster IgG (H&L) Antibody Fluorescein Conjugated Pre-Adsorbed
<b>Applications:</b>	IF
<b>Recommended Dilution:</b>	<b>IF:</b> 1:500-1:2500 <b>FLISA:</b> 1:10,000-1:50,000
<b>Host:</b>	Goat
<b>Immunogen:</b>	Golden Syrian and Armenian Hamster IgG whole molecules
<b>Formulation:</b>	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
<b>Reconstitution Method:</b>	Restore with deionized water (or equivalent) - Reconstitution Volume: 500 µL
<b>Concentration:</b>	1.0 mg/mL - lot specific
<b>Conjugation:</b>	FITC
<b>Storage:</b>	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
<b>Note:</b>	F(ab') <sub>2</sub> Anti-Golden Syrian & Armenian Hamster IgG Fluorescein Antibody has been tested by western blot and is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.



[View online »](#)

**Product images:**



Western blot of Fluorescein conjugated Goat F(ab')<sub>2</sub> Anti-Hamster IgG Pre-Adsorbed secondary antibody. Lane 1: Golden Syrian Hamster IgG. Lane 2: Armenian Hamster IgG. Load: 50 ng per lane. Primary antibody: None. Secondary antibody: Fluorescein goat secondary antibody at 1:1,000 for 60 min at RT. Blocking: MB-070 for 30 min at RT. Predicted/Observed size: 25 & 55 kDa, 25 & 55 kDa for Hamster IgG. Other band(s): None.