

## **Product datasheet for TA398345**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Fab Sheep IgG (H&L) Antibody Peroxidase Conjugated

**Product data:** 

**Product Type:** Secondary Antibodies

**Product Name:** Fab Sheep IgG (H&L) Antibody Peroxidase Conjugated

**Applications:** ELISA, IHC, WB

Recommended Dilution: WB: 1:1,000 - 1:10,000

IHC: 1:500 - 1:2,500

**ELISA**: 1:10,000 - 1:50,000

Host: Rabbi

Immunogen: Sheep IgG whole molecule

**Formulation:** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

**Reconstitution Method:** Restore with deionized water (or equivalent) - Reconstitution Volume: 500 μL

**Concentration:** 1.0 mg/mL - lot specific

Conjugation: HRP

Storage: Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -

20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as

an undiluted liquid. Dilute only prior to immediate use.

Note: Fab Anti-Sheep IgG Peroxidase Antibody has been tested by ELISA and is suitable for


immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot

consistency, high titer and specificity.





## **Product images:**



ELISA Results of Fab Rabbit Anti-Sheep IgG Antibody Peroxidase Conjugated. Each well was coated in duplicate with 1.0  $\mu$ g of Sheep IgG (p/n 013-0102). The starting dilution of antibody was 5 $\mu$ g/ml and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using TMB substrate (p/n TMBE-1000).