

Product datasheet for TA400025

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

BIRC5 Mouse Monoclonal Antibody [Clone ID: OTI2D9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2D9

Applications: WB

Recommended Dilution: WB: 1:1000

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human BIRC5 (NP_001159) produced in HEK293T

cell

Formulation: PBS (pH7.4) containing 50% glycerol, 0.1% BSA

Concentration: 0.5 mg/ml

Purification: Purified from cell culture supernatant by affinity chromatography

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 16.2 kDa

Gene Name: baculoviral IAP repeat containing 5

Database Link: NP 001159

Entrez Gene 332 Human

015392



Background:


This gene is a member of the inhibitor of apoptosis (IAP) gene family, which encode negative regulatory proteins that prevent apoptotic cell death. IAP family members usually contain multiple baculovirus IAP repeat (BIR) domains, but this gene encodes proteins with only a single BIR domain. The encoded proteins also lack a C-terminus RING finger domain. Gene expression is high during fetal development and in most tumors yet low in adult tissues. Antisense transcripts are involved in the regulation of this gene's expression. At least four transcript variants encoding distinct isoforms have been found for this gene, but the full-length natures of only three of them have been determined.

Synonyms: API4; EPR-1

Protein Families: Druggable Genome, Stem cell - Pluripotency

Protein Pathways: Colorectal cancer, Pathways in cancer

Product images:



HEK293T cells were transfected with pCMV6-ENTRY control (Lane 1) or pCMV6-ENTRY BIRC5 (Lane 2) plasmids for 48 hrs. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BIRC5-HRP.