

Product datasheet for **TA400028**

Isocitrate dehydrogenase (IDH1) Mouse Monoclonal Antibody [Clone ID: OTI2H9]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI2H9
Applications:	FC
Recommended Dilution:	FC: 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full-length protein expressed in 293T cell transfected with human IDH1 expression vector
Formulation:	PBS (pH7.4) containing 50% glycerol, 0.1% BSA and 0.02% NaN ₃
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids by affinity chromatography
Conjugation:	DyLight 488
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	46.7 kDa
Gene Name:	isocitrate dehydrogenase (NADP(+)) 1
Database Link:	NP_005887 Entrez Gene 3417 Human O75874



[View online »](#)

Background:


Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. Each NADP(+)-dependent isozyme is a homodimer. The protein encoded by this gene is the NADP(+)-dependent isocitrate dehydrogenase found in the cytoplasm and peroxisomes. It contains the PTS-1 peroxisomal targeting signal sequence. The presence of this enzyme in peroxisomes suggests roles in the regeneration of NADPH for intraperoxisomal reductions, such as the conversion of 2, 4-dienoyl-CoAs to 3-enoyl-CoAs, as well as in peroxisomal reactions that consume 2-oxoglutarate, namely the alpha-hydroxylation of phytanic acid. The cytoplasmic enzyme serves a significant role in cytoplasmic NADPH production.

Synonyms:

HEL-216; HEL-S-26; IDCD; IDH; IDP; IDPC; PICD

Protein Pathways:

Citrate cycle (TCA cycle), Glutathione metabolism, Metabolic pathways

Product images:

HEK293T cells were transfected with pCMV6-ENTRY control (PS01, left panel) or IDH1 cDNA (right panel) and incubated for 48 hours after transfection. The cells were detached, fixed with 2% formaldehyde for 30 min, permeabilized with 0.1% triton-X100, and then stained with PE-labeled IDH1 antibodies (TA400028).