

## **Product datasheet for TA420102**

## Phospho-Chk2 (CHEK2) Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB,1:1000 - 1:5000

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

**Immunogen:** A synthetic phosphorylated peptide around T68 of human Chk2 (NP\_009125.1).

**Formulation:** Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Stable for 12 months from date of receipt.

**Predicted Protein Size:** 61kDa - Observed MW: 62kDa

**Gene Name:** checkpoint kinase 2

**Database Link:** Entrez Gene 11200 Human


<u>096017</u>



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

In response to DNA damage and replication blocks, cell cycle progression is halted through the control of critical cell cycle regulators. The protein encoded by this gene is a cell cycle checkpoint regulator and putative tumor suppressor. It contains a forkhead-associated protein interaction domain essential for activation in response to DNA damage and is rapidly phosphorylated in response to replication blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. In addition, this protein interacts with and phosphorylates BRCA1, allowing BRCA1 to restore survival after DNA damage. Mutations in this gene have been linked with Li-Fraumeni syndrome, a highly penetrant familial cancer phenotype usually associated with inherited mutations in TP53. Also, mutations in this gene are thought to confer a predisposition to sarcomas, breast cancer, and brain tumors. This nuclear protein is a member of the CDS1 subfamily of serine/threonine protein kinases. Several transcript variants encoding different isoforms have been found for this gene.

Synonyms:

CDS1; CHK2; hCds1; HuCds1; LFS2; PP1425; RAD53