

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA421198

CL2A Rabbit Monoclonal Antibody [Clone ID: 1G9]

Product data:

Product Type:	Primary Antibodies
Clone Name:	1G9
Applications:	ELISA
Recommended Dilution:	ELISA 1:5000-10000
Host:	Rabbit
lsotype:	IgG
Clonality:	Monoclonal
Reconstitution Method:	Reconstitute with sterile, low-endotoxin water. See the COA for more details
Concentration:	Lot specific
Purification:	Purified from cell culture supernatant by affinity chromatography
Conjugation:	Unconjugated
Storage:	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing)
Stability:	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background:	CL2A is a crucial component in antibody-drug conjugate (ADC) therapy, acting as a chemical linker that connects a potent DNA topoisomerase I inhibitor, SN-38, to an antibody. Representing Cysteine-Linked 2-Aminoethyl, CL2A features a complex structure with a PEG8 chain, a triazole ring, a PABC-peptide, and a maleimide group. The maleimide group binds to a cysteine residue on the antibody, facilitating targeted drug delivery. Designed to release SN- 38 in acidic cancer cell environments, CL2A induces DNA damage and cell death. This versatile linker is employed in ADCs targeting various antigens like Trop-2 or HER2, tailored to specific cancer types.



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Product images:




Figure 1. Elisa plates were pre-coated with IgG-CL2A (0.2µg/per well). Serial diluted anti-CL2A monoclonal antibody (TA421198) solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-CL2A monoclonal antibody binding with IgG-CL2A is 12.57ng/ml.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US