

Product datasheet for TA421199AM

CL2A Rabbit Monoclonal Antibody [Clone ID: 1H2]

Product data:

Product Type: Primary Antibodies

Clone Name: 1H2
Applications: ELISA

Recommended Dilution: ELISA 1:5000-10000

Host: Rabbit Isotype: IgG

Clonality: Monoclonal

Reconstitution Method: Reconstitute with sterile, low-endotoxin water. See the COA for more details

Concentration: Lot specific

Purification: Purified from cell culture supernatant by affinity chromatography

Conjugation: Biotin

Storage: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended

for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing)

Stability: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended

for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.

Background: CL2A is a crucial component in antibody-drug conjugate (ADC) therapy, acting as a chemical

linker that connects a potent DNA topoisomerase I inhibitor, SN-38, to an antibody.

Representing Cysteine-Linked 2-Aminoethyl, CL2A features a complex structure with a PEG8 chain, a triazole ring, a PABC-peptide, and a maleimide group. The maleimide group binds to a cysteine residue on the antibody, facilitating targeted drug delivery. Designed to release SN-38 in acidic cancer cell environments, CL2A induces DNA damage and cell death. This versatile linker is employed in ADCs targeting various antigens like Trop-2 or HER2, tailored to specific

cancer types.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:

ELISA assay to evaluate Anti-CL2A Antibody 0.2µg Human IgG-CL2A per well




Figure 1. Elisa plates were pre-coated with IgG-CL2A (0.2µg/per well). Serial diluted anti-CL2A monoclonal antibody ([TA421199]) solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-CL2A monoclonal antibody binding with IgG-CL2A is 2.801ng/ml.