

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA421205

# Eribulin Rabbit Monoclonal Antibody [Clone ID: 3E2]

## **Product data:**

Product Type:	Primary Antibodies
Clone Name:	3E2
Applications:	ELISA
Recommended Dilution:	ELISA 1:5000-10000
Host:	Rabbit
lsotype:	lgG
Clonality:	Monoclonal
Reconstitution Method:	Reconstitute with sterile, low-endotoxin water. See the COA for more details
Concentration:	Lot specific
Purification:	Purified from cell culture supernatant by affinity chromatography
Conjugation:	Unconjugated
Storage:	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing)
Stability:	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background:	Eribulin, a fully synthetic macrocyclic ketone analogue derived from the marine sponge Halichondria okadai, shows promise as an ADC payload. Its anti-tumor activity involves inhibiting microtubule elongation and mitotic spindle formation, leading to apoptosis. Eribulin-based ADCs offer advantages such as well-understood toxicity profiles and compatibility with various linker chemistries. Clinical development includes MORAb-202, an ADC targeting folate receptor alpha (FRα) in solid tumors.



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



## **Product images:**




Figure 1. Elisa plates were pre-coated with IgG-Eribulin (0.2µg/per well). Serial diluted anti-Eribulin monoclonal antibody (TA421205) solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Eribulin monoclonal antibody binding with IgG-Eribulin is 2.587ng/ml.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US