

Product datasheet for TA421208S

DM1 Rabbit Monoclonal Antibody [Clone ID: 14E3]

Product data:

Product Type: Primary Antibodies

Clone Name: 14E3
Applications: ELISA

Recommended Dilution: ELISA 1:5000-10000

Host: Rabbit Isotype: IgG

Clonality: Monoclonal

Reconstitution Method: Reconstitute with sterile, low-endotoxin water. See the COA for more details

Concentration: Lot specific

Purification: Purified from cell culture supernatant by affinity chromatography

Conjugation: Unconjugated

Storage: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended

for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing)

Stability: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended

for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.

Background: DM1, also known as mertansine, represents a promising class of ADCs. As a microtubulin

inhibitor, DM1 disrupts microtubule assembly by binding to tubulin at the rhizoxin binding site. The ingenious concept behind ADCs involves coupling DM1 to a monoclonal antibody, enabling targeted delivery to tumor cells. Notably, trastuzumab emtansine (T-DM1) has already gained approval for HER2-positive breast cancer treatment, exemplifying the

potential of DM1-based ADCs in precision oncology.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:

ELISA assay to evaluate Anti-DM1 Antibody 0.2µg BSA-DM1 per well




Figure 1. Elisa plates were pre-coated with BSA-DM1 (0.2µg/per well). Serial diluted anti-DM1 monoclonal antibody ([TA421208]) solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-DM1 monoclonal antibody binding with BSA-DM1 is 2.110ng/ml.