

Product datasheet for TA426152

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

EU: info-de@origene.com CN: techsupport@origene.cn

IKZF1 Rabbit Monoclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, WB

Recommended Dilution: WB,1:500 - 1:2000

IF/ICC,1:50 - 1:200

ELISA,Recommended starting concentration is 1 μg/mL. Please optimize the concentration

based on your specific assay requirements. ChIP,5µg antibody for 10µg-15µg of Chromatin

CUT&Tag, 10⁵ cells /1 µg

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Monoclonal

Formulation: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 58kDa





Background:

This gene encodes a transcription factor that belongs to the family of zinc-finger DNA-binding proteins associated with chromatin remodeling. The expression of this protein is restricted to the fetal and adult hemo-lymphopoietic system, and it functions as a regulator of lymphocyte differentiation. Several alternatively spliced transcript variants encoding different isoforms have been described for this gene. Most isoforms share a common C-terminal domain, which contains two zinc finger motifs that are required for hetero- or homo-dimerization, and for interactions with other proteins. The isoforms, however, differ in the number of N-terminal zinc finger motifs that bind DNA and in nuclear localization signal presence, resulting in members with and without DNA-binding properties. Only a few isoforms contain the requisite three or more N-terminal zinc motifs that confer high affinity binding to a specific core DNA sequence element in the promoters of target genes. The non-DNA-binding isoforms are largely found in the cytoplasm, and are thought to function as dominant-negative factors. Overexpression of some dominant-negative isoforms have been associated with B-cell malignancies, such as acute lymphoblastic leukemia (ALL).