

## **Product datasheet for TA426532**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **DDX17 Rabbit Monoclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

**Applications:** ELISA, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC-P,1:50 - 1:200

ELISA, Recommended starting concentration is 1 µg/mL. Please optimize the concentration

based on your specific assay requirements.

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Monoclonal

Formulation: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 80kDa

**Background:** DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are

putative RNA helicases. They are implicated in a number of cellular processes involving

alteration of RNA secondary structure, such as translation initiation, nuclear and

mitochondrial splicing, and ribosome and splicesosome assembly. Based on their distribution

patterns, some members of this family are believed to be involved in embryogenesis,

spermatogenesis, and cellular growth and division. This gene encodes a DEAD box protein, which is an ATPase activated by a variety of RNA species, but not by dsDNA. This protein, and

that encoded by DDX5 gene, are more closely related to each other than to any other member of the DEAD box family. This gene can encode multiple isoforms due to both

alternative splicing and the use of alternative translation initiation codons, including a non-AUG (CUG) start codon.

