

Product datasheet for **TA500023S**

GSC Mouse Monoclonal Antibody [Clone ID: OTI1D7]


Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI1D7
Applications:	FC, IHC, WB
Recommended Dilution:	WB 1:500~1000, IHC 1:50, FLOW 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 107-257 of human GSC (NP_776248) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.7 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	28.0 kDa
Gene Name:	goosecoid homeobox
Database Link:	NP_776248 Entrez Gene 145258 Human P56915
Background:	Goosecoid is a member of the bicoid subfamily of the paired (PRD) homeobox family of proteins. It acts as a transcription factor and may be autoregulatory. A similar protein in mice plays a role in craniofacial and rib cage development during embryogenesis.
Synonyms:	SAMS
Protein Families:	Transcription Factors




[View online »](#)


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY GSC (Cat# [RC209586], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-GSC(Cat# [TA500023]). Positive lysates [LY406285] (100ug) and [LC406285] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-GSC monoclonal antibody.




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-GSC mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500023])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human kidney tissue using anti-GSC mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500023])




Immunohistochemical staining of paraffin-embedded Human liver tissue within the normal limits using anti-GSC mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500023])



Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-GSC mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500023])



Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-GSC mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500023])



HEK293T cells transfected with either [RC209586] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-GSC antibody ([TA500023]), and then analyzed by flow cytometry.